

Table 6A.3.1.1: Trends in Deaths Due to Falls, by Age, United States 2000-2010

Year	Total Unintentional Injury Deaths Occurring All Locations			Age-Adjusted Rate per 100,000 Persons [1]
	<u>Total</u>	<u>Death</u>	<u>% of Deaths</u>	
	<u>Unintentional</u>	<u>due to Falls</u>	<u>due to Falls</u>	
	<u>Injury Deaths</u>			
2000	97,900	13,300	13.6%	4.82
2001	101,500	15,000	14.8%	5.33
2002	106,700	16,300	15.3%	5.69
2003	109,300	17,200	15.7%	5.93
2004	112,000	18,800	16.8%	6.37
2005	117,800	19,700	16.7%	6.53
2006	121,600	20,800	17.1%	6.77
2007	123,700	22,600	18.3%	7.21
2008	121,900	24,000	19.7%	7.50
2009	118,000	24,800	21.0%	7.58
2010	120,900	26,000	21.5%	7.82
2000-2010	1,251,400	218,600	17.5%	6.56

	% of Unintentional Injury Deaths Due to Falls by Age					
	<u>≤24</u>	<u>25-34</u>	<u>35-44</u>	<u>45-54</u>	<u>55-64</u>	<u>65+</u>
2000	1.6%	2.6%	3.9%	7.1%	12.6%	33.1%
2001	1.9%	2.9%	4.1%	7.7%	13.1%	35.6%
2002	1.7%	2.4%	4.0%	6.9%	13.0%	38.2%
2003	1.7%	2.3%	3.8%	6.6%	13.3%	39.9%
2004	1.7%	2.5%	4.0%	7.0%	14.4%	42.5%
2005	1.5%	2.1%	3.6%	6.4%	13.4%	43.0%
2006	1.6%	2.1%	3.6%	6.6%	13.9%	45.4%
2007	1.6%	2.2%	3.5%	6.4%	14.3%	47.9%
2008	1.7%	2.0%	3.4%	6.4%	14.2%	50.2%
2009	1.7%	2.1%	3.6%	6.7%	14.6%	52.2%
2010	1.7%	2.1%	3.3%	6.5%	14.3%	52.4%

[1] Age-adjusted to 2000 population, all races, both sexes.

Source: Centers for Disease Control. Web-based Injury Statistics Query and Reporting System (WISQARS) [online]. (2011) National Center for Injury Prevention and Control (producer). Available from URL: <http://www.cdc.gov/injury/wisqars/fatal.html>. [October 24, 2013].