Table 1.2.2: Prevalence and Age-Adjusted Rate of Self-Reported Select Medical Conditions by Age,United States 2012

							Prevalence Per 100 Persons				
	Total Persons With Condition Aged 18 & Over (in millions)					in Age Group					Rate [9] Per
Medical Condition	<u>18-44</u>	<u>45-64</u>	65-74	<u>75+</u>	Total	18-44	45-64	65-74	<u>75+</u>	Total	100 Total
Musculoskeletal [1]	46.188	51.156	16.655	12.648	126.647	41.6	62.4	70.1	69.9	53.9	52.8
Emotional Disorder [2]	41.224	31.177	6.439	4.720	83.560	37.1	38.0	27.1	26.1	35.6	34.8
Circulatory [3]	12.658	32.104	14.376	12.987	72.125	11.4	39.1	60.5	71.8	30.7	30.0
Sleep Disorder	27.849	26.059	7.910	6.628	68.446	25.1	31.8	33.3	36.6	29.1	28.5
Respiratory [4]	26.851	26.355	7.808	5.118	66.132	24.2	32.1	32.9	28.3	28.2	27.6
Migraines or Severe Headaches	24.847	15.181	2.514	1.410	43.952	22.4	18.5	10.6	7.8	18.7	18.3
Mental Health Disorder [5]	20.055	15.823	3.910	2.227	42.015	18.1	19.3	16.5	12.3	17.9	17.5
Diabetes	5.326	14.647	6.827	4.435	31.234	4.8	17.9	28.7	24.5	13.3	13.0
Blind or Trouble Seeing w/Glasses	6.014	9.292	2.607	2.696	20.609	5.4	11.3	11.0	14.9	8.8	8.6
Cancer [6]	2.265	7.629	5.014	5.165	20.073	2.0	9.3	21.1	28.6	8.5	8.4
Cognitive Disorder [7]	4.799	7.185	2.328	3.435	17.747	4.3	8.8	9.8	19.0	7.6	7.4
Hearing Trouble [8]	1.890	5.140	3.108	4.998	15.137	1.7	6.3	13.1	27.6	6.4	6.3
Kidney Disease	0.633	1.548	0.746	0.954	3.882	0.6	1.9	3.1	5.3	1.7	1.6
Liver Disease	0.688	1.662	0.491	0.193	3.034	0.6	2.0	2.1	1.1	1.3	1.3

[1] Includes arthritis, chronic joint symptoms (more than 3 months in duration), pain in lower back or neck or radiating leg pain (more than 3 months duration).

[2] Includes feelings that interfere with life and activities in past 30 days: sad, nervous, restless, hopelessness, worthlessness, depression, general anxiety, panic disorder, and living is an effort.

[3] Includes coronary heart disease, angina pectoris, heart attack, stroke, chronic hypertension, heart disease.

[4] Includes emphysema, chronic asthma, hay fever, sinusitis, chronic bronchitis.

[5] Includes ADD/ADHD, bipolar disorder, depression in past 12 months, or other mental health disorder .

[6] Includes all types of cancer.

[7] Includes memory loss or neurological problem in past 12 months.

[8] Includes respondents reporting "moderate trouble," "a lot of trouble" or "deaf" without a hearing aid or hearing device, or currently uses a hearing aid.

[9] Age-adjusted by direct method to US Census population estimate for July 1, 2012. Accessed October 26, 2013. NOTE: Due to sample weighting to approximate the age distribution of the U.S. population, the crude (unadjusted) and age-adjusted rates of reported conditions are very close.

Source: National Health Interview Survey (NHIS)_Adult sample. www.cdc.gov/nchs/nhis/nhis_2012_data_release.htm. July 2, 2013.