

**Table 5.3:** Percentages of Individuals Hospitalized or Admitted to Skilled Nursing Facility within Six Months After Various Fracture Types Compared With Six Months Preceding the Fracture, United States 2000-2004

	Site of Fracture						
	<u>Clinical</u> <u>Vertebral</u>	<u>Hip</u>	<u>Pelvis</u>	<u>Femur</u>	<u>Tibia/</u> <u>Fibula</u>	<u>Humerus</u>	<u>Distal</u> <u>Radius/</u>
<b>Hospital Discharges</b>							
% Hospitalized six months after fracture	58.4	98.2	91.4	90.5	70.0	62.9	35.4
% Hospitalized in baseline (six months before)	41.5	27.1	34.2	40.5	36.8	36.2	28.5
<b>Skilled Nursing Facility Stays After Fracture</b>							
% Stays after fracture	34.7	84.1	72.2	75.3	49.8	38.4	19.2
% Stays in baseline (six months before)	26.3	24.0	30.0	32.3	30.1	26.8	16.7

Source: Becker DJ, Yun H, Kilgore ML, et al. Health services utilization after fractures: recent evidence from Medicare. *J Gerontol A Biol Sci Med Sci*. 2010 Sep;65(9):1012-20. PMID: 20530242