

**Table 6A.2.1: Patient Visits for Injuries as Proportion of All Health Care Visits, by Sex and Age, United States 2010/2011**

	Total Visits (in 000s) [7]	Proportion of All Patient Visits							
		All Visits	Sex		Age in Years				
			Male	Female	<18	18-44	45-64	65-74	75 & over
Proportion of General Population			49.2%	50.8%	24.0%	36.5%	26.4%	7.0%	6.0%
<b>HOSPITAL VISITS</b>									
NIS 2011 [1]									
All Patient Visits	38,590.7	100.0%	42.0%	58.0%	14.7%	24.3%	25.1%	13.9%	21.9%
Visits for Musculoskeletal Injury	1,708.6	4.4%	44.1%	55.9%	4.6%	18.2%	23.9%	13.9%	39.5%
Visits for Other Injuries	797.7	2.1%	58.4%	41.6%	6.8%	23.4%	26.0%	12.7%	31.1%
NHDS 2010 [2]									
All Patient Visits	38,919.2	100.0%	41.7%	58.3%	16.1%	24.6%	24.4%	13.3%	21.6%
Visits for Musculoskeletal Injury	1,693.9	4.4%	42.9%	57.1%	7.1%	18.5%	24.3%	12.4%	37.7%
Visits for Other Injuries	784.3	2.0%	59.1%	40.9%	8.3%	22.1%	24.9%	12.9%	31.8%
<b>EMERGENCY DEPARTMENTS</b>									
NEDS 2010 [3]									
All Patient Visits	128,970.4	100.0%	44.5%	55.5%	19.8%	40.6%	22.6%	6.9%	10.2%
Visits for Musculoskeletal Injury	19,010.2	14.7%	51.0%	49.0%	22.8%	41.0%	20.9%	5.6%	9.7%
Visits for Other Injuries	7,630.1	5.9%	54.1%	45.9%	27.9%	41.6%	18.1%	4.3%	8.1%
NHAMCS_ED 2010 [4]									
All Patient Visits	129,843.4	100.0%	44.9%	55.1%	23.0%	40.6%	21.5%	6.3%	8.7%
Visits for Musculoskeletal Injury	19,421.6	15.0%	52.3%	47.7%	26.3%	42.0%	18.3%	4.9%	8.5%
Visits for Other Injuries	7,620.2	5.9%	57.2%	42.8%	32.9%	39.8%	16.3%	4.3%	6.6%
<b>OUTPATIENT CLINICS</b>									
NHAMCS_OP 2010 [5]									
All Patient Visits	100,742.1	100.0%	40.4%	59.6%	24.7%	28.0%	27.5%	10.6%	9.2%
Visits for Musculoskeletal Injury	3,856.2	3.8%	52.3%	47.7%	5.6%	3.8%	3.2%	2.1%	3.0%
Visits for Other Injuries	1,010.6	1.0%	53.5%	46.5%	1.9%	1.0%	0.6%	0.3%	0.6%
<b>PHYSICIAN OFFICES</b>									
NAMCS 2010 [6]									
All Patient Visits	1,008,802.0	100.0%	41.8%	58.2%	19.0%	26.0%	29.4%	13.1%	12.6%
Visits for Musculoskeletal Injury	40,799.1	4.0%	52.7%	47.3%	4.1%	4.9%	4.3%	2.8%	3.0%
Visits for Other Injuries	9,740.8	1.0%	50.4%	49.6%	0.6%	1.5%	1.1%	0.5%	0.6%

[1] Source: HCUP Nationwide Inpatient Sample (NIS). Healthcare Cost and Utilization Project (HCUP). 2011. Agency for Healthcare Research and Quality, Rockville, MD. [www.hcup-us.ahrq.gov/nisoverview.jsp](http://www.hcup-us.ahrq.gov/nisoverview.jsp)

[2] Source: National Hospital Discharge Survey (NHDS), 2010. [www.cdc.gov/nchs/nhds/nhds\\_questionnaires.htm](http://www.cdc.gov/nchs/nhds/nhds_questionnaires.htm) April 23, 2013.

[3] Source: HCUP Nationwide Emergency Department Sample (NEDS). Healthcare Cost and Utilization Project (HCUP). 2010. Agency for Healthcare Research and Quality, Rockville, MD. [www.hcup-us.ahrq.gov/nedsoverview.jsp](http://www.hcup-us.ahrq.gov/nedsoverview.jsp)

[4] Source: National Hospital Ambulatory Medical Care Survey\_Emergency Department (NHAMCS\_ED), 2010. [www.cdc.gov/nchs/ahcd/ahcd\\_questionnaires.htm](http://www.cdc.gov/nchs/ahcd/ahcd_questionnaires.htm) April 23, 2013.

[5] Source: National Hospital Ambulatory Medical Care Survey\_Outpatient Department (NHAMCS\_OP), 2010. [www.cdc.gov/nchs/ahcd/ahcd\\_questionnaires.htm](http://www.cdc.gov/nchs/ahcd/ahcd_questionnaires.htm) April 23, 2013.

[6] Source: National Ambulatory Medical Care Survey (NAMCS), 2010. [www.cdc.gov/nchs/ahcd/ahcd\\_questionnaires.htm](http://www.cdc.gov/nchs/ahcd/ahcd_questionnaires.htm) April 23, 2013.

[7] Total visits may be lower than sum of diagnoses due to multiple diagnoses per patient.