

Table T7B.5.2: Estimated Number of US Residents with Osteoporosis and Low Bone Mass in 2010 by Sex and State

State	Population (in 100,000s)								
	Male			Female			Both Sexes		
	Total	With Condition		Total	With Condition		Total	With Condition	
		Osteoporosis	Low Bone Mass		Osteoporosis	Low Bone Mass		Osteoporosis	Low Bone Mass
Alabama	7.28	0.31	2.56	8.65	1.33	4.45	15.93	1.64	6.99
Alaska	1.02	0.04	0.36	0.95	0.15	0.49	1.97	0.2	0.87
Arizona	9.5	0.41	3.34	10.74	1.65	5.52	20.24	2.08	8.88
Arkansas	4.51	0.19	1.59	5.21	0.8	2.68	9.73	1	4.27
California	50.63	2.18	17.82	57.83	8.91	29.72	108.46	11.17	47.61
Colorado	7.22	0.31	2.54	7.96	1.23	4.09	15.18	1.56	6.67
Connecticut	5.65	0.24	1.99	6.69	1.03	3.44	12.34	1.27	5.42
Delaware	1.4	0.06	0.49	1.66	0.26	0.85	3.06	0.32	1.34
District of Columbia	0.75	0.03	0.26	0.95	0.15	0.49	1.7	0.18	0.75
Florida	32.03	1.38	11.28	37.34	5.75	19.19	69.38	7.15	30.46
Georgia	12.69	0.55	4.47	15.01	2.31	7.72	27.7	2.85	12.16
Hawaii	2.22	0.1	0.78	2.47	0.38	1.27	4.69	0.48	2.06
Idaho	2.31	0.1	0.81	2.48	0.38	1.28	4.8	0.49	2.11
Illinois	18.44	0.79	6.49	21.69	3.34	11.15	40.13	4.13	17.62
Indiana	9.66	0.42	3.4	11.17	1.72	5.74	20.83	2.15	9.14
Iowa	4.9	0.21	1.73	5.58	0.86	2.87	10.49	1.08	4.6
Kansas	4.27	0.18	1.5	4.85	0.75	2.49	9.12	0.94	4
Kentucky	6.65	0.29	2.34	7.71	1.19	3.96	14.37	1.48	6.31
Louisiana	6.57	0.28	2.31	7.66	1.18	3.94	14.23	1.47	6.25
Maine	2.41	0.1	0.85	2.73	0.42	1.4	5.14	0.53	2.26
Maryland	8.4	0.36	2.96	10.04	1.55	5.16	18.44	1.9	8.1
Massachusetts	10.03	0.43	3.53	12	1.85	6.17	22.03	2.27	9.67
Michigan	15.72	0.68	5.53	18.07	2.78	9.29	33.79	3.48	14.83
Minnesota	8.11	0.35	2.85	9.03	1.39	4.64	17.14	1.77	7.53
Mississippi	4.27	0.18	1.5	5.09	0.78	2.62	9.36	0.96	4.11
Missouri	9.27	0.4	3.26	10.79	1.66	5.55	20.05	2.07	8.8
Montana	1.77	0.08	0.62	1.88	0.29	0.96	3.64	0.38	1.6
Nebraska	2.77	0.12	0.97	3.13	0.48	1.61	5.9	0.61	2.59
Nevada	4	0.17	1.41	4.22	0.65	2.17	8.23	0.85	3.61
New Hampshire	2.23	0.1	0.78	2.46	0.38	1.26	4.69	0.48	2.06
New Jersey	13.21	0.57	4.65	15.86	2.44	8.15	29.07	2.99	12.76
New Mexico	3.18	0.14	1.12	3.59	0.55	1.84	6.76	0.7	2.97
New York	28.65	1.23	10.08	34.77	5.35	17.87	63.42	6.53	27.84
North Carolina	13.88	0.6	4.89	16.55	2.55	8.51	30.43	3.13	13.36
North Dakota	1.1	0.05	0.39	1.2	0.18	0.62	2.3	0.24	1.01
Ohio	18.24	0.78	6.42	21.37	3.29	10.99	39.61	4.08	17.39
Oklahoma	5.64	0.24	1.99	6.47	1	3.33	12.12	1.25	5.32
Oregon	6.21	0.27	2.19	6.98	1.08	3.59	13.19	1.36	5.79
Pennsylvania	20.94	0.9	7.37	24.73	3.81	12.71	45.66	4.7	20.05
Rhode Island	1.64	0.07	0.58	1.99	0.31	1.02	3.64	0.37	1.6
South Carolina	7.06	0.3	2.48	8.36	1.29	4.3	15.42	1.59	6.77
South Dakota	1.3	0.06	0.46	1.44	0.22	0.74	2.74	0.28	1.2
Tennessee	9.65	0.41	3.4	11.34	1.75	5.83	20.99	2.16	9.21
Texas	32.21	1.38	11.34	36.54	5.63	18.78	68.74	7.08	30.18
Utah	3.07	0.13	1.08	3.35	0.52	1.72	6.42	0.66	2.82
Vermont	1.11	0.05	0.39	1.23	0.19	0.63	2.34	0.24	1.03
Virginia	11.69	0.5	4.11	13.56	2.09	6.97	25.25	2.6	11.08
Washington	10.24	0.44	3.6	11.34	1.75	5.83	21.58	2.22	9.47
West Virginia	3.31	0.14	1.17	3.74	0.58	1.92	7.05	0.73	3.1
Wisconsin	9.04	0.39	3.18	10.09	1.55	5.19	19.13	1.97	8.4
Wyoming	0.93	0.04	0.33	0.95	0.15	0.49	1.88	0.19	0.82
United States	458.97	19.74	161.56	531.51	81.85	273.2	990.49	102.02	434.82

Wright NC, Looker AC, Saag KG, et al. The recent prevalence of osteoporosis and low bone mass in the United States based on bone mineral density at the femoral neck or lumbar spine. J Bone Miner Res. 2014;29(11):2520-2526. doi:10.1002/jbmr.2269.