

Table 5F.1.2.3: Hospitalizations for Injury/Poisoning Events¹, Active Category, US Armed Forces 2008-2017 by Sex

Year	Hospitalizations			Share of Total [5]	
	Total [2]	Males	Females	Males	Females
2008	10,282	9,201	1,081	89.5%	10.5%
2009	11,879	9,937	1,219	89.1%	10.9%
2010	11,855	9,791	1,159	89.4%	10.6%
2011	12,095	11,410	1,331	89.6%	10.4%
2012	10,684	9,986	1,154	89.6%	10.4%
2013	8,599	7,348	970	88.3%	11.7%
2014	7,415	6,364	963	86.9%	13.1%
2015	7,034	5,969	881	87.1%	12.9%
2016	6,752	5,822	930	86.2%	13.8%
2017	6,467	5,603	864	86.6%	13.4%

[1] Burden of disease major categories and burden of disease-related conditions based on a modified version of those defined in the Global Burden of Disease Study (Source: The global burden of disease: a comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projected to 2020. Murray CJ, Lopez AD, eds. Harvard School of Public Health (on behalf of the World Health Organization and The World Bank), 1996:120–122.

[2] Total hospitalizations, rate of hospitalization an ambulatory encounters values changed across the reporting years when three (alternate) years were reported. Figures shown are based on the most recent summary reports included in multi-year tables rather than the primary reporting year summary.

[3] Percentages adjusted to original reported number of total encounters as subtables were not refined in subsequent years.

Source: Armed Forces Health Surveillance Branch. Absolute and relative morbidity burdens attributable to various illnesses and injuries, active component, US Armed Forces, 2017. *MSMR* Annual Summary Edition, 2009 thru 2018.