

**Table 5F.1.3.4: Most Frequent Diagnoses During Ambulatory Visits with ICD-10 Codes, by Major Diagnostic Category, by Sex, US Armed Forces, 2015, 2016, and 2017**

	2015				2016				2017			
	Females		Males		Females		Males		Females		Males	
	Number	%	Number	%	Number	%	Number	%	Number	%	Number	%
Sprain of ankle (and foot)	10,678	10.6%	46,085	9.4%	11,736	8.3%	48,162	6.9%	11,234	8.3%	45,439	7.1%
Sprains and strains of knee (cruciate ligament) and leg	10,630	10.6%	44,610	9.1%	4,146	2.9%	26,039	3.7%	4,603	3.4%	25,672	4.0%
Sprains and strains of other and unspecified parts of back	7,000	7.0%	28,082	5.7%								
Injury other and unspecified	6,545	6.5%	34,521	7.1%								
Certain adverse effects not elsewhere classified	6,177	6.2%										
Sprains and strains of shoulder and upper arm			36,747	7.5%	2,989	2.1%	30,967	4.4%			27,293	4.3%
Sprain of hip					34,373	2.4%			3,178	2.4%		
Concussion									3,038	2.3%	15,097	2.4%
Injury of other muscles and tendons at lower leg level					5,262	3.7%			2,287	2.1%		
Fracture of other and unspecified metacarpal bone							15,271	2.2%			15,199	2.4%
History of TBI (traumatic brain injury)							17,378	2.5%				

Source: Armed Forces Health Surveillance Branch. Ambulatory Visits, Active Component, U.S. Armed Forces, (year). TABLE 2 (3). Most frequent diagnoses during ambulatory visits with ICD-10 codes, by major diagnostic category, males (females), active component, US Armed Forces, (year). *MSMR Annual Summary Edition*, 20016, 2017, 2018.