

Table 1.4.4: Prevalence of Chronic Joint Pain [1] for Persons Age 18 and Over, by Joint, by Geographic Region, United States 2015

<u>Joint Pain Site</u>	<u>Total Persons With Condition Aged 18 & Over (in Millions)</u>					<u>Prevalence Per 100 Persons in Regional Group</u>				
	<u>Northeast</u>	<u>Midwest</u>	<u>South</u>	<u>West</u>	<u>Total</u>	<u>Northeast</u>	<u>Midwest</u>	<u>South</u>	<u>West</u>	<u>Total</u>
Knee	8.132	11.283	17.974	9.753	47.142	18.2	21.3	19.5	16.9	19.0
Shoulder	3.934	4.906	8.638	4.850	22.327	8.8	9.3	9.4	8.4	9.0
Hip	2.867	4.523	7.123	3.665	18.177	6.4	8.5	7.7	6.3	7.3
Fingers	2.676	4.640	6.550	4.245	18.112	6.0	8.7	7.1	7.3	7.3
Ankle	2.240	3.315	6.037	3.184	14.776	5.0	6.3	6.5	5.5	6.0
Wrist	2.001	3.089	5.263	3.142	13.492	4.5	5.8	5.7	5.4	5.4
Elbow	1.749	2.711	4.315	2.286	11.061	3.9	5.1	4.7	4.0	4.5
Toes	0.971	1.828	3.210	1.557	7.565	6.5	8.8	3.5	2.7	3.1
Other Joint	0.868	0.884	1.534	1.095	4.381	1.9	1.7	1.7	1.9	1.8
All Chronic Joint [2]	12.233	17.447	26.761	16.091	72.532	27.4	32.9	29.0	27.9	29.3
Mean Chronic Joint Sites (1 to 9)	2.1	2.1	2.3	2.1	2.2	4.7	4.0	2.5	3.6	0.9

* Does not meet standards for reliability.

[1] Symptoms lasting 3 months or longer

[2] Chronic pain in multiple joints may be reported

Source: National Health Interview Survey (NHIS)_Adult sample, 2015. <http://www.cdc.gov/nchs/nhis/data-questionnaires-documentation.htm> July 23, 2016.