

Table 3A.2.1.1: Site of Joint Pain for Persons with Self-Reported Doctor-Diagnosed Arthritis, Adults Age 18 and Older, by Sex, United States 2013-2015

Site of Joint Pain [1]	Adults with Self-Reported Doctor-Diagnosed Arthritis (DDA) [2] (in 1,000s)					
	Male	Rate Per 100 DDA Population	Female	Rate Per 100 DDA Population	Total	Rate Per 100 DDA Population
Knee	9,866.4	44.7	15,673.7	48.5	25,540.1	47.0
Shoulder	5,570.0	25.3	7,347.9	22.7	12,917.9	23.8
Finger	4,011.1	18.2	7,810.0	24.2	11,821.1	21.7
Hip	3,904.6	17.7	7,738.8	23.9	11,643.4	21.4
Ankle	3,344.2	15.2	5,567.6	17.2	8,911.8	16.4
Wrist	2,847.0	12.9	5,479.7	17.0	8,326.8	15.3
Elbow	2,696.4	12.2	3,791.8	11.7	6,668.2	12.3
Toes	1,942.6	8.8	3,399.8	10.5	5,342.4	9.8
Other	911.4	4.1	1,531.5	4.7	2,442.9	4.5
All Sites Joint Pain	15,916.6	72.2	24,277.7	75.1	40,194.3	73.9

[1] Self-reported joint pain lasting 30 days or longer. Joint pain is self-reported and is not equivalent to medical code defined arthropathic joint pain included in AORC tables.

[2] Responded "yes" when asked: Have you ever been told by a doctor or other health professional that you have some form of arthritis, rheumatoid arthritis, gout, lupus, or fibromyalgia?

Source: National Health Interview Survey (NHIS)_Adult sample, 2013, 2014, 2015. <http://www.cdc.gov/nchs/nhis/data-questionnaires-documentation.htm> July 23, 2016.