

**Table 3A.2.1.5:** Proportion of Multiple Sites and Mean Number of Self-Reported Joint Pain Sites for Adults Age 18 & Older with Reported DDA and Chronic Joint Pain, United States, 2013-2015

Number of Joint Pain Sites Reported [1]	Proportion of Adults with Joint Pain in at Least One Site													
	All Adults	Males	Females	18-44	45-64	65+	Non-Hispanic Whites	Non-Hispanic Blacks	Hispanic	Non-Hispanic Other	Northeast	Midwest	South	West
1	40.5%	43.1%	38.7%	41.7%	36.9%	44.1%	40.0%	44.8%	38.1%	41.8%	44.9%	39.1%	40.2%	38.8%
2	25.6%	26.4%	25.1%	24.3%	26.1%	25.4%	25.9%	24.5%	24.0%	26.5%	25.9%	25.5%	25.6%	25.5%
3	14.2%	13.0%	14.6%	13.7%	15.1%	12.8%	13.9%	14.6%	15.5%	11.7%	14.6%	13.6%	13.8%	14.3%
4	6.6%	5.8%	7.2%	6.7%	7.1%	6.0%	6.9%	4.8%	7.1%	6.0%	5.5%	7.1%	6.6%	7.0%
5	3.8%	3.1%	4.2%	3.3%	4.1%	3.6%	3.8%	3.3%	3.8%	4.3%	2.8%	4.2%	3.7%	4.3%
6	2.6%	2.7%	2.5%	2.4%	3.2%	2.0%	2.7%	2.0%	2.6%	2.5%	1.7%	2.7%	2.7%	2.9%
7	2.1%	1.7%	2.3%	2.1%	2.5%	1.5%	1.9%	2.2%	2.9%	3.1%	1.0%	2.4%	2.2%	2.3%
8	4.3%	3.7%	4.6%	4.9%	4.4%	3.9%	4.2%	3.7%	5.5%	3.9%	2.8%	4.7%	4.6%	4.4%
9	0.7%	0.5%	0.8%	0.7%	0.7%	0.6%	0.8%	0.2%	0.5%	0.1%	0.8%	0.6%	0.7%	0.5%
Mean sites reported	2.5	2.4	2.6	2.5	2.6	2.4	2.5	2.3	2.6	2.4	2.2	2.6	2.5	2.6
Number of Adults with DDA Reporting Joint Pain (in 000s)	40,194.3	15,916.6	24,277.7	5,475.3	18,446.6	16,272.4	30,513.7	4,627.2	3,210.7	1,842.7	7,064.1	9,970.8	15,048.7	8,110.7
Proportion of DDA with Chronic Joint Pain	73.9%	72.2%	75.1%	68.7%	76.3%	73.2%	73.8%	76.0%	72.5%	73.1%	71.2%	74.4%	74.4%	74.8%

[1] Self-reported joint pain lasting 30 days or longer. Sites include knee, shoulder, finger, hip, ankle, wrist, elbow, toes, other. Joint pain is self-reported and is not equivalent to medical code defined arthropathic joint pain included in AORC tables. Joint pain can be caused by arthritis, injuries, or degeneration of bone surfaces.

Source: National Health Interview Survey (NHIS)\_Adult sample, 2013, 2014, 2015. <http://www.cdc.gov/nchs/nhis/data-questionnaires-documentation.htm> July 23, 2016.