

Table 4A.1.1: Prevalence of Osteoporosis and Low Bone Mass at the Femur Neck and Lumbar Spine, Adults Aged 50+ Years, United States 2013-2014

	Number	Femur Neck		Lumbar Spine		Either Femur Neck or Lumbar Spine	
		Percent	SE [1]	Percent	SE [1]	Percent	SE [1]
Osteoporosis [2]							
Both Sexes	2001	6.3%	0.6	7.8%	0.4	11.0%	0.7
Female	1029	9.8%	1.1	11.6%	1	16.5%	1.2
Male	972	2.5%	0.5	3.6%	0.7	5.1%	0.7
<i>p</i> value, sex		<0.001			<0.001		<0.001
Low Bone Mass [3]							
Both Sexes	2001	42.8%	1.0	27.9%	1.5	44.5%	1.0
Female	1029	52.6%	1.5	35.9%	1.7	52.6%	1.1
Male	972	32.1%	1.3	19.2%	1.8	35.6%	1.3
<i>p</i> value, sex		<0.001			<0.001		<0.001

[1] Standard error.

[2] Osteoporosis was defined as a T-score \leq -2.5.

[3] Low bone mass was defined as a T-Score between -1.0 and -2.5.

Source: Looker AC, Sarafrazi Isfajani N, Fan B, Shepherd JA. Trends in osteoporosis and low bone mass in older US adults, 2005-2006 through 2013-2014. *Osteoporos Int*, 2017. Jun;28(6):1979-1988. doi: 10.1007/s00198-017-3996-1. Reproduced with permission.