

**Table 4A.2.8:** Trend in Age-adjusted Hip Fracture Incidence Rate Among Women in Medicare, United States 2010-2015

	Age-adjusted Hip Fracture Incidence Rate per 1,000 Person-Years				5% Sample Confidence Interval		
	<u>5% Sample</u>		<u>Postmenopausal Osteoporosis Cohort</u>		<u>Person Years</u>	<u>Crude Incidence Rate</u>	<u>Age-standardized Incidence Rate</u>
2010	7517	8.05	49571	13.68	798872	9.41 (9.20, 9.62)	8.05 (7.87, 8.24)
2011	7999	8.65	54094	14.42	794240	10.07 (9.85, 10.29)	8.65 (8.46, 8.85)
2012	7708	8.42	54604	14.33	781431	9.86 (9.64, 10.08)	8.42 (8.22, 8.61)
2013	7700	8.60	58953	14.16	772569	9.97 (9.74, 10.19)	8.60 (8.40, 8.80)
2014	7419	8.52	61089	14.61	765632	9.69 (9.47, 9.91)	8.52 (8.33, 8.72)
2015	5846	8.30	58890	14.93	675302	8.66 (8.43, 8.88)	8.30 (8.09, 8.52)
Change 2015-2010		3.1%		9.1%			
Change 2015-2012		-1.4%		4.2%			

Source: Lewiecki ME, Wright NC, Curtis JR, et al., Hip fracture trends in the United States, 2002 to 2015. *Osteoporos Int*, 2018. 29(3): p. 717-722.